Mesenteric approach during pancreaticoduodenectomy for pancreatic ductal adenocarcinoma






MAPLE-PD trial

Department of Surgery, Wakayama Medical University

Hiroki Yamaue






- (1) Posterior approach
 Ohigashi et al. *Hepatogastroenterology 2004*Pessaux et al. *J Gastrointest Surg 2006*
- (2) Medical unicinate approach Shukla et al. *Hepatogastroenterology 2007*
- (3) CLIP method Kawai and Yamaue. World J Surg 2008
- (4) Left posterior approach Horiguchi et al. *JHBPS 2007* Kurosaki et al. *JOP 2011*
- (5) Mesenteric approach
 Nakao et al. *Hepatogastroenterology 1993*Weitz et al. *J Am Coll Surg 2010*



^{&#}x27;Artery-first' approaches to pancreatoduodenecotmy. Sanjay et al. *Br J Surg 2012;99:1027-1935*



(4) Left posterior approach Horiguchi et al. *JHBPS 2007* Kurosaki et al. *JOP 2011*

(5) Mesenteric approach
Nakao et al. *Hepatogastroenterology 1993*Weitz et al. *J Am Coll Surg 2010*

Kawai and Yamaue. World J Surg 2008

'Artery-first' approaches to pancreatoduodenecotmy. Sanjay et al. *Br J Surg 2012;99:1027-1935*





(1) Posterior approach
Ohigashi et al. *Hepatogastroenterology 2004*Pessaux et al. *J Gastrointest Surg 2006*

"How I Do It"

Pancreaticoduodenectomy: Superior Mesenteric Artery First Approach

Patrick Pessaux, M.D., Deepak Varma, M.D., Jean-Pierre Arnaud, M.D.




Kurosaki et al. *JOP 2011*

(5) Mesenteric approach Nakao et al. *Hepatogastroenterology 1993* Weitz et al. *J Am Coll Surg 2010*

'Artery-first' approaches to pancreatoduodenecotmy. Sanjay et al. *Br J Surg 2012;99:1027-1935*






- (1) Posterior approach
 Ohigashi et al. *Hepatogastroenterology 2004*Pessaux et al. *J Gastrointest Surg 2006*
- (2) Medical unicinate approach Shukla et al. *Hepatogastroenterology 2007*
- (3) CLIP method Kawai and Yamaue. World J Surg 2008
- (4) Left posterior approach Horiguchi et al. *JHBPS 2007* Kurosaki et al. *JOP 2011*
- (5) Mesenteric approach
 Nakao et al. *Hepatogastroenterology 1993*Weitz et al. *J Am Coll Surg 2010*

'Artery-first' approaches to pancreatoduodenecotmy. Sanjay et al. *Br J Surg 2012;99:1027-1935*





- (1) Posterior approach Ohigashi et al. *Hepatogastroenterology 2004* Pessaux et al. *J Gastrointest Surg 2006*
- (2) Medical unicinate approach Shukla et al. *Hepatogastroenterology 2007*
- (3) CLIP method

SURGEON AT WORK

The "Artery First" Approach for Resection of Pancreatic Head Cancer

Jürgen Weitz, MD, Nuh Rahbari, MD, Moritz Koch, MD, Markus W Büchler, MD


IVI

Weitz et al. J Am Coll Surg 2010

'Artery-first' approaches to pancreatoduodenecotmy. Sanjay et al. *Br J Surg 2012;99:1027-1935*



Wakayama Medical university



- (1) Posterior approach
 Ohigashi et al. *Hepatogastroenterology 2004*Pessaux et al. *J Gastrointest Surg 2006*
- (2) Medical unicinate approach
 Shukla et al. *Hepatogastroenterology 2007*
- (3) CLIP method Kawai and Yamaue. World J Surg 2008
- (4) Left posterior approach Horiguchi et al. *JHBPS 2007* Kurosaki et al. *JOP 2011*

(5) Mesenteric approach Nakao et al. Hepatogastroenterology 1993

Weitz et al. JAm Coll Surg 2010



(1) Posterior approach
Ohigashi et al. *Hepatogastroenterology 2004*Pessaux et al. *J Gastrointest Surg 2006*

Isolated Pancreatectomy for Pancreatic Head Carcinoma Using Catheter Bypass of the Portal Vein

A. Nakao, H. Takagi

Department of Surgery II, Nagoya University School of Medicine, Nagoya, Japan



(5) Mesenteric approach Nakao et al. Hepatogastroenterology 1993

Weitz et al. JAm Coll Surg 2010

Concept of artery-first approach during PD for PDAC

Artery-first approach

PD starts from dissection of connective tissues around the SMA and division of the inferior pancreaticoduodenal artery (IPDA)


The aims of artery-first approach

- ✓ Early determination of the resectability status for PDAC during PD
- ✓ Control of pancreatic head congestion by early division of the IPDA
 - → Decrease of intraoperative blood loss volume
- **✓** Complete dissection around the SMA
 - → Increase of the R0 rate

Posterior approach as artery-first approach


Posterior approach: exposing the origin of superior mesenteric artery (SMA) in front of the left renal vein after kocherization.

	author	publish	year	number	page
1	Ohigashi et al.	Hepatogastroenterology	2004	51	4-5
2	Pessaux et al.	J Gastrointest Surg	2006	10	921-926
3	Traian et al.	Langenbecks Arch Surg	2010	395	677-684



Medial uncinate approach as artery-first approach


Medial uncinate approach: Retrograde resection of the pancreatic head is performed starting with the uncinate process after division of the first jejunal loop and transection of the pancreas as the last operative step of the resection.




Left posterior approach as artery-first approach

Left posterior approach: exposing J1 and J2 arteries at their origin on the SMA in the transverse mesocolon. As next step, division of the IPDA arising from the posterior surface of the SMA.


author		publish	year number		page	
1	Horiguchi et al.	JHBPS	2007	14	575-578	
2	Kurosaki et al.	JOP	2011	12	220-229	





Kurosaki et al. JOP 2011; 12: 220-229


The methods of artery-first approaches



Mesenteric approach as artery-first approach


First, exposing the SMA-SMV at the base of the transverse mesocolon


	author	publish	year	number	page
1	Nakao et al.	Hepatogastroenterology	1993	40	426-429
2	Weitz et al.	J Am Coll Surg	2010	210	e1-e4
3	Nakao et al.	Dig Surg	2016	33	308-13
4	Hirono et al.	Ann Gastroenterol Surg	2017	in press	



Area of incision of the peritoneum at duodenojejunal flexure (dotted line).

Mesenteric approach





Mesentric approach can facilitate dissection of posterior soft tissues of SMA and SMV, so called mesoduodenum and mesopancreas, at the first step of Whipple's operation.

Wakayama Medical university

Pancreaticoduodenectomy with CHA resection (PD-CHAR)

Patient; 50-y, female

Diagnosis; Pancreas Neck ~ Body Cancer with common hepatic artery

and portal vein invasion

NAC; GEM + nab-paclitaxel 2 courses → NAC-GA trial

(NCT02926183, UMIN000024154)

Effects of NAC; Tumor diameter 30mm → 30mm (SD)

PET-CT SUVmax3.38 \rightarrow 2.07

CA19-9 88.9 \rightarrow 66.0 U/ml, CEA 1.3 \rightarrow 1.7 ng/ml

Operation; PD-CHAR


Operation time; 8hr 27min, Bleeding 585 ml, No blood tranfusion

Pathology; CH(+) DU(-) S(+) RP(+) PV(+) Ach(+) Plcha(+) OO(-) n(+:8a)

T4(PV,A) N1 M0 StageIVa



Preoperative MD-CT findings; CHA invasion




Preoperative MD-CT findings; SMV Portal vein invasion









T4 pancreatic cancer in Neck and Body




Splenic artery was removed 6cm Taped the root of CHA




Splenic artery and PHA anastomosis






Reconstruction of portal vein



Pancreaticoduodenectomy with CHA resection



Operation; PD-CHAR, Arterial anastomosis under microscope

Op time; 8hr 27min

Bleeding; 585 ml

Clinical benefits of mesenteric approach

We evaluated the clinical benefits of mesenteric approach, especially survival benefits during pancreaticoduodenectomy (PD) for pancreatic ductal adenocarcinoma (PDAC) patients.

This is a <u>first report regarding the long-term survival</u>* of Mesenteric approach.

* Hirono S, Yamaue H, et al. Mesenteric approach during pancreaticoduodenectomy for pancreatic ductal adenocarcinoma.

Ann Gastroenterol Surg 2017, in press



Patients and methods

[Study design] Retrospective cohort study

[Patients] 237 PD patients 2000 to 2015 at WMUH

• Conventional approach group: 165 patients (2000~2011)

• Mesenteric approach group: 72 patients (2011~2015)

[Surgical procedure]

<u>Conventional approach</u>: During resection procedure, after Kocher's maneuver at first step, regional lymph node dissection and resection of stomach, bile duct, and pancreas, and then, <u>dissection of connective tissue around the SMA</u>, including <u>diving inferior pancreaticoduodenal artery (IPDA) at the final step</u>.

Mesenteric approach: Dissection connected tissue around the SMA, including diving IPDA was performed at the first step.

Statistical analysis

Matched-pairs analysis

: propensity score approach using a multiple logistic regression model with 4 covariates: <u>age</u>, <u>sex</u>, <u>resectability status</u>, and <u>neoadjuvant therapy</u>



• Conventional approach: 58 patients

• Mesenteric approach: 58 patients

(Resectable PDAC: 58 patients, BR PDAC: 58 patients)



Backgrounds: mesenteric vs. conventional

	Resectable I	PDAC	BR PDAC	
	Mesenteric (n=28)	Convention (n=30)	Mesenteric (n=30)	Convention (n=28)
Age, median	67(42-82)	69 (43-87)	70.5 (48-78)	70 (49-79)
Sex, male	18 (64%)	22 (73%)	15 (50%)	17 (61%)
Serum CA19-9 U/ml	135	140	525	349
Resectability (NC	CN)			
Resectable	28 (100%)	30 (100%)	0	0
BR	0	0	30 (100%)	28 (100%)
BR-AV			10 (33%)	9 (32%)
BR-A			8 (27%)	9 (32%)
BR-V			12 (40%)	10 (36%)
Tumor size, mm	23.6	22.0	28.7	27.3
Neoadjuvant therapy	0	0	9 (30%)	9 (32%)
Follow-up (mo)	22.5	27.6	11.7	12.0

Surgical outcome for resectable PDAC Mesenteric vs. conventional approach

	Mesenteric approach	Conventional approach	
	(n=28)	(n=30)	P
PV resection, n (%)	11 (39%)	6 (20%)	0.107
Operative time, min,	416	371	0.007
median (range)	(314-535)	(254-	520)
Blood loss volume, ml,	313	502	0.023
median (range)	(40-1500)	(60-2230)	
Transfusion, n (%)	1 (4%)	6 (20%)	0.055
Mobility, n (%)	5 (18%)	3 (10%)	0.386
Mortality, n (%)	0	1 (3%)	0.330
Hospital stay, day,	14	21	0.007
median (range)	(10-36)	(11-65)	

Surgical outcome for BR PDAC Mesenteric vs. conventional approach

	Mesenteric approach (n=30)	Conventional approach (n=28)	P
PV resection, n (%)	24 (80%)	21 (75%)	0.648
Operative time, min, median (range)	459 (374-620)	452 (322-570)	0.210
Blood loss volume, ml, median (range)	507 (115-2225)	920 (115-3610)	0.003
Transfusion, n (%)	5 (17%)	13 (46%)	0.014
Mobility, n (%)	4 (13%)	5 (18%)	0.634
Mortality, n (%)	0	0	-
Hospital stay, day, median (range)	15 (9-59)	17.5 (10-42)	0.863


Pathological findings for resectable PDAC Mesenteric vs. conventional approach

	Mesenteric approach (n=28)	Conventional approach (n=30)	P
Number of metastatic lymph nodes			0.229
0	7 (25%)	14 (47%)	
1-3	16 (57%)	12 (40%)	
≧4	5 (18%)	4 (13%)	
Number of harvested lymph nodes,	23	23.5	0.919
n, median (range)	(11-53)	(11-48)	
R0 rate	28 (100%)	26 (87%)	0.045
Adjuvant therapy within 8 weeks after surgery, n (%)	19 (68%)	17 (57%)	0.380
Completion of planned postoperative adjuvant therapy	20 (71%)	16 (53%)	0.156

Pathological findings for BR PDAC Mesenteric vs. conventional approach

	Mesenteric approach	Conventional approach	
	(n=30)	(n=28)	\boldsymbol{P}
Number of metastatic lymph nodes			0.114
0	6 (20%)	6 (21%)	
1-3	9 (30%)	15 (54%)	
≧4	15 (50%)	7 (25%)	
Number of harvested lymph nodes,	26.5	26	0.668
n, median (range)	(10-53)	(9-49)	
R0 rate	24 (80%)	24 (86%)	0.565
Adjuvant therapy within 8 weeks after surgery, n (%)	18 (60%)	14 (50%)	0.444
Completion of planned postoperative adjuvant therapy	12 (40%)	10 (36%)	0.737


Overall survival for resectable PDAC Mesenteric vs. conventional approach



MST
Mesenteric (-)
Conventional 18.7 m



Overall survival for BR PDAC Mesenteric vs. conventional approach



MST
Mesenteric approach 11.6 m
Conventional approach 13.3 m

BR



Summary and conclusion

- ✓ The blood loss volume and transfusion frequency were significantly lower in mesenteric approach group than in conventional approach group for both resectable PDAC and BR PDAC.
- ✓ The R0 rate was significantly higher in mesenteric approach group than in conventional approach group for resectable PDAC.
- → Mesenteric approach has survival benefits for resectable PC.
- ✓ There were no significant differences of R0 rate and overall survival between mesenteric and conventional approach groups for BR PDAC.
- \rightarrow Mesenteric approach might not have oncological benefit for BR PDAC, and multidisciplinary treatments are essential to improve the survival for BR PDAC patients.

MAPLE-PD trial

Mesenteric APproach vs. conventionaL approach for pancrEatic cancer during PD

Multicenter Randomized Clinical Trial:

Primary endpoint: Overall survival

Secondary endpoints: Operative time, Blood loss,

Transfusion, Complications,

R0 rate, Disease-free survival

Accompanying research:

Circulating tumor DNA in portal vein blood during operation

We plan to start this RCT on August 2017!

